

Epimorphisms and Boundary Slopes of 2-Bridge Knots

JIM HOSTE
PATRICK D. SHANAHAN

In this article we study a partial ordering on knots in S^3 where $K_1 \geq K_2$ if there is an epimorphism from the knot group of K_1 onto the knot group of K_2 which preserves peripheral structure. If K_1 is a 2-bridge knot and $K_1 \geq K_2$, then it is known that K_2 must also be 2-bridge. Furthermore, Ohtsuki, Riley, and Sakuma give a construction which, for a given 2-bridge knot $K_{p/q}$, produces infinitely 2-bridge knots $K_{p'/q'}$ with $K_{p'/q'} \geq K_{p/q}$. After characterizing all 2-bridge knots with 4 or less distinct boundary slopes, we use this to prove that in any such pair, $K_{p'/q'}$ is either a torus knot or has 5 or more distinct boundary slopes. We also prove that 2-bridge knots with exactly 3 distinct boundary slopes are minimal with respect to the partial ordering. This result provides some evidence for the conjecture that all pairs of 2-bridge knots with $K_{p'/q'} \geq K_{p/q}$ arise from the Ohtsuki-Riley-Sakuma construction.

57M25

1 Introduction

Many interesting problems in knot theory involve the question of when one knot complement can be mapped onto another, or at the algebraic level, when the fundamental group of one knot complement can be mapped onto the fundamental group of another. For example, Simon's Conjecture asserts that the fundamental group of any knot in S^3 can surject onto only finitely many knot groups. This conjecture has been established for 2-bridge knots by Boileau, Boyer, Reid and Wang [2]. On the other hand, given any 2-bridge knot $K_{p/q}$, Ohtsuki, Riley, and Sakuma [16] show how to construct infinitely many 2-bridge knots $K_{p'/q'}$ whose complements map onto the complement of $K_{p/q}$. Their construction induces an epimorphism $\phi : \pi_1(S^3 - K_{p'/q'}) \rightarrow \pi_1(S^3 - K_{p/q})$ of knot groups which *preserves peripheral structure*, that is, ϕ takes the subgroup generated by a longitude and meridian of $K_{p'/q'}$ into a conjugate of the subgroup generated by a longitude and meridian of $K_{p/q}$.

For any pair of knots, not just 2-bridge knots, Silver and Whitten define a partial ordering $\geq$ on knots in S^3 by declaring that $K_1 \geq K_2$ if there exists an epimorphism

$\phi : \pi_1(S^3 - K_1) \rightarrow \pi_1(S^3 - K_2)$ which preserves peripheral structure [18]. Thus, for a given 2-bridge knot $K_{p/q}$, the Ohtsuki-Riley-Sakuma (ORS) construction gives infinitely many 2-bridge knots $K_{p'/q'}$ such that $K_{p'/q'} \geq K_{p/q}$. This leads naturally to the following question posed in [16].

Question 1.1 (*Ohtsuki-Riley-Sakuma*) *Is every pair of 2-bridge knots $(K_{p'/q'}, K_{p/q})$ with $K_{p'/q'} \geq K_{p/q}$ given by the ORS construction?*

González-Acuña and Ramírez [6] give an affirmative answer to Question 1.1 in the case where $K_{p/q}$ is a 2-bridge torus knot. In this paper we give additional evidence that the answer to Question 1.1 is yes in general. Numerous computations with non-torus knot ORS pairs $(K_{p'/q'}, K_{p/q})$ with $K_{p'/q'} \geq K_{p/q}$ suggest that in any such pair the larger knot $K_{p'/q'}$ must be sufficiently complex with respect to a variety of knot invariants such as: crossing number, degree of the Alexander polynomial, degree of the A-polynomial, number of distinct boundary slopes, etc. Thus, if the answer to Question 1.1 is yes, then knots with “small” complexity should be minimal. Here, a knot K_1 is said to be *minimal* with respect to the Silver–Whitten partial ordering if $K_1 \geq K_2$ implies that either $K_2 = K_1$, K_2 is the mirror image of K_1 , or K_2 is the unknot. Throughout the remainder of this paper, we will consider two knots K_1 and K_2 to be equivalent, $K_1 \equiv K_2$, if either $K_2 = K_1$ or K_2 is the mirror image of K_1 .

One of the themes of this paper will be to look at the set of distinct boundary slopes of a knot. These are easily computed for 2-bridge knots due to the classification of Hatcher and Thurston [7]. Our first main result gives a lower bound on the number of distinct boundary slopes for the larger knot in an ORS pair.

Theorem 4.2 *If $K_{p'/q'} \geq K_{p/q}$ is a nontrivial ORS pair, then either*

- i. *$K_{p'/q'}$ and $K_{p/q}$ are both torus knots and $K_{p'/q'}$ has precisely two distinct boundary slopes, or*
- ii. *$K_{p'/q'}$ has at least five distinct boundary slopes.*

Our second main result establishes the minimality of 2-bridge knots with exactly three boundary slopes.

Theorem 5.1 *If $K_{p/q}$ is a 2-bridge knot with exactly 3 distinct boundary slopes, then $K_{p/q}$ is minimal with respect to the Silver–Whitten partial order.*

Notice that if the answer to Question 1.1 is yes, then Theorem 4.2 implies Theorem 5.1. Thus, we may think of Theorem 5.1 as evidence in support of an affirmative answer to Question 1.1. It is also worth noting that Theorem 4.2 is sharp. That is, there exist non-minimal 2-bridge knots with 5 distinct slopes. We provide an example of this in Section 4.

This project began as joint work of the first author and two undergraduate students, Tómasz Przytycki and Rebecca Nachison, in the summer of 2007 at the Claremont Colleges REU program. Thanks are due to Tómasz and Rebecca for working out most of Theorem 3.2. We thank the National Science Foundation, the Claremont Colleges, and Pitzer College in particular, for their support of the REU program. The second author would also like to thank the Claremont Colleges for their hospitality and Pitzer College for its support during the completion of this work. Finally, thanks are due to Alan Reid for helpful conversations regarding character varieties.

2 Preliminaries

We begin with some notation for 2-bridge knots. Recall that each 2-bridge knot corresponds to a relatively prime pair of integers p and q with q odd. We denote the knot as $K_{p/q}$. Furthermore, $K_{p/q}$ and $K_{p'/q'}$ are ambient isotopic as unoriented knots if and only if $q' = q$ and $p' \equiv p^{\pm 1} \pmod{q}$ (see [4] for details). As mentioned earlier, we will not distinguish between a knot $K_{p/q}$ and its mirror image $K_{-p/q}$. Therefore, in this paper we consider two 2-bridge knots $K_{p/q}$ and $K_{p'/q'}$ to be equivalent if and only if $q' = q$ and either $p' \equiv p^{\pm 1} \pmod{q}$ or $p' \equiv -p^{\pm 1} \pmod{q}$. Because of this classification, it is sometimes convenient to assume that $0 < p < q$.

The theory of 2-bridge knots is closely tied to continued fractions. We adopt the convention used in [16] and define $p/q = r + [b_1, b_2, \dots, b_n]$ as the continued fraction

$$p/q = r + [b_1, b_2, \dots, b_n] = r + \frac{1}{b_1 + \frac{1}{b_2 + \frac{\ddots}{\ddots + \frac{1}{b_n}}}}$$

It is well known that if $0 < p < q$ then we may write p/q uniquely as $p/q = 0 + [b_1, b_2, \dots, b_n]$ where each $b_i > 0$ and $b_n > 1$. Furthermore, we may also assume




Figure 1: A 4-plat diagram for $K_{p/q}$ where $p/q = r + [b_1, b_2, \dots, b_n]$ (n even).

that $b_1 > 1$ for the following reason. If $b_1 = 1$, reversing the order of the partial quotients will give $p'/q = [b_n, \dots, b_3, b_2, 1] = [b_n, \dots, b_3, b_2 + 1]$ with $0 < p' < q$ and $pp' \equiv (-1)^{n+1} \pmod{q}$. Thus, for any 2-bridge knot K , there exists a continued fraction $0 + [b_1, b_2, \dots, b_n]$ with $b_i > 0$, $b_1 > 1$, and $b_n > 1$ representing K (or its mirror image). In what follows, we will refer to such a continued fraction as *strongly positive*. Moreover, we will also refer to a vector $(b_1, b_2, \dots, b_n)$ of positive integers with $b_1 > 1$ and $b_n > 1$ as *strongly positive*. Finally, the negation of such a vector we call *strongly negative*.

If $p/q = r + [b_1, b_2, \dots, b_n]$, then $K_{p/q}$ can be represented by the standard 4-plat diagram shown in Figure 1. In each box there are b_i crossings. If i is odd, then the crossings are right-handed if $b_i > 0$ and left-handed otherwise. For i even the convention reverses, with $b_i > 0$ corresponding to left-handed crossings and $b_i < 0$ for right-handed. Thus, the 4-plat diagram is alternating precisely when all b_i have the same sign. Notice also that there are two possible ways to “close up” the strands at the right end of the braid in Figure 1 depending on whether n is even or odd. Figure 1 depicts a 4-plat with n even. If n is odd, strings 1 and 2, and strings 3 and 4, connect pairwise at the right end of the braid just as they do at the left end. With this notation, the fraction $1/3 = [3]$ gives the right-hand trefoil knot while $2/5 = [2, 2]$ gives the figure eight knot.

Now suppose that ϕ is a homomorphism between knot groups that preserves peripheral structure. Since all nontrivial knots in S^3 have Property P [12], there are certain restrictions placed on the images of a meridian and longitude of K_1 .

Theorem 2.1 *Let K_1 and K_2 be nontrivial knots with $K_1 \geq K_2$. Then there exists an epimorphism $\phi : \pi_1(S^3 - K_1) \rightarrow \pi_1(S^3 - K_2)$ and an integer d such that*

$$(1) \quad \phi(\mu_1) = \mu_2, \text{ and}$$

$$(2) \quad \phi(\lambda_1) = \lambda_2^d$$

for some choice $\{\mu_i, \lambda_i\}$ of meridian-(preferred) longitude pairs for K_i .

Proof. Since ϕ preserves peripheral structure we may assume (after conjugation, if necessary) that

$$\begin{aligned}\phi(\mu_1) &= \mu_2^a \lambda_2^b, \text{ and} \\ \phi(\lambda_1) &= \mu_2^c \lambda_2^d\end{aligned}$$

for some integers a, b, c and d . If we consider the abelianization map $ab : \pi_1(S^3 - K_2) \rightarrow \mathbb{Z}$, then $ab(\phi(\lambda_1)) = 0$ since λ_1 is in the commutator subgroup. However, $ab(\phi(\lambda_1)) = ab(\mu_2^c \lambda_2^d) = c$. This shows that $c = 0$. Moreover, $ab(\phi(\mu_1)) = ab(\mu_2^a \lambda_2^b) = a$ must generate $\mathbb{Z}$ because $ab \circ \phi$ is onto and $\pi_1(S^3 - K_1)$ is generated by conjugates of μ_1 . Therefore, $a = \pm 1$. By replacing μ_2 with its inverse if necessary, we may assume that $a = 1$.

Now consider $1/b$ Dehn filling on K_2 . This filling kills the image of μ_1 and any of its conjugates, and so kills the entire group, since ϕ is onto. Thus the manifold we obtain after filling is simply connected. Since K_2 is a nontrivial knot with Property P, we must have done trivial surgery. Therefore, $b = 0$. $\square$

Because of Theorem 2.1, it makes sense to write

$$K_1 \geq_d K_2$$

whenever $K_1 \geq K_2$ and there exists an epimorphism ϕ as in Theorem 2.1 with $\phi(\lambda_1) = \lambda_2^d$. In general, d can take on any integer value. In particular, given the knot K_2 and any integer d , there exists a knot K_1 and epimorphism ϕ realizing d . The reader is referred to [11] for entré into this subject. On the other hand, if K_1 and K_2 are 2-bridge knots, then d must be odd.

Theorem 2.2 *If $K_{p'/q'} \geq_d K_{p/q}$ where $K_{p'/q'}$ and $K_{p/q}$ are 2-bridge knots and $K_{p/q}$ is nontrivial, then d is odd. In particular, the image $\phi(\lambda_1)$ under the epimorphism ϕ cannot be trivial.*

Proof. Suppose $\phi : \pi_1(S^3 - K_{p'/q'}) \rightarrow \pi_1(S^3 - K_{p/q})$ is an epimorphism taking λ_1 to λ_2^d and let ρ be an irreducible parabolic representation of $\pi_1(S^3 - K_{p/q})$ into $SL(2, \mathbb{C})$ (such a representation exists by Riley [17]). Composing ρ with ϕ gives such a representation for $K_{p'/q'}$. It follows from [8] that any irreducible parabolic representation of a 2-bridge knot into $SL(2, \mathbb{C})$ must take the longitude into the conjugacy class of an upper triangular matrix with diagonal entries of -1 . Hence ϕ cannot take λ_1 to an even power of λ_2 . $\square$

The following theorem summarizes some additional properties of the partial ordering which we will refer to in this article. The first two parts are contained in [18], the third comes from [2], and the fourth appears in [1].

Theorem 2.3 (Silver-Whitten, Boileau-Boyer, Boileau-Boyer-Reid-Wang) Suppose that $K_1 \geq_d K_2$.

- i. If $\Delta_K^{(i)}(t)$ is the i -th Alexander polynomial of K , then $\Delta_{K_2}^{(i)}(t)$ divides $\Delta_{K_1}^{(i)}(t)$.
- ii. If $A_K(L, M)$ is the A-polynomial of K , then $A_{K_2}(L, M)$ divides $(L^d - 1)A_{K_1}(L^d, M)$.
- iii. If $X(K)$ is the $SL(2, \mathbb{C})$ character variety of K , then the induced map $\phi^* : X(K_2) \rightarrow X(K_1)$ is an injective, algebraic and closed (in the Zariski topology) mapping.
- iv. If K_1 is a 2-bridge knot, then K_2 is a 2-bridge knot. Furthermore, if $K_1 = K_{p'/q'}$ and $K_2 = K_{p/q}$, then either $K_1 \equiv K_2$ or $q' = kq$ with $k > 1$.

Notice that the fact that q divides q' in Theorem 2.3(iv) is easily derived from (i) and the fact that $q = |\Delta_{K_{p/q}}(-1)|$ is the determinant of $K_{p/q}$.

Because of the relationship between the A-polynomial and boundary slopes of a knot, Theorem 2.3(ii) enables us to use boundary slopes as a tool to study whether one knot is greater than another. Recall that a *boundary slope* $r = \frac{a}{b}$ of a knot K is an element of $\mathbb{Q} \cup \infty$ such that K contains a properly embedded, incompressible, boundary incompressible surface S in its exterior whose boundary is the curve $a\mu + b\lambda$ (or multiple copies of this curve). Here $\{\mu, \lambda\}$ is a meridian-(preferred) longitude pair for K . The *Newton polygon* of the A-polynomial $A_K(L, M) = \sum a_{ij}L^iM^j$ is the convex hull of the set of points $\{(i, j) \in \mathbb{R}^2 \mid a_{ij} \neq 0\}$. If r is the slope of a side of the Newton polygon of $A_K(L, M)$, then it is proven in [5], that r is a boundary slope of K . A boundary slope of K which appears as a slope of the Newton polygon is called *strongly detected*. We shall tacitly assume that the A-polynomial always includes the component $L - 1$ corresponding to abelian representations so that 0 is always a strongly detected slope.

Corollary 2.4 Suppose $K_1 \geq_d K_2$, $d \neq 0$, and that r is a strongly detected boundary slope of K_2 . Then dr is a strongly detected boundary slope of K_1 .

Proof. We establish this result using the following well-known property of Newton polygons. For any polynomial $P(x, y)$ let S_P denote the set of slopes of the sides of the Newton polygon of P . Then for all polynomials $P(x, y)$ and $Q(x, y)$, $S_P \cup S_Q = S_{PQ}$ (a proof may be found in [8]). Now suppose that r is a strongly detected boundary slope of K_2 . Then r is the slope of a side of the Newton polygon of $A_{K_2}(L, M)$. Since $A_{K_2}(L, M)$ divides $(L^d - 1)A_{K_1}(L^d, M)$ it follows that $r = 0$ or r is a slope of a side of the Newton polygon of $A_{K_1}(L^d, M)$. If $r = 0$, then $dr = 0$ is a strongly detected slope of K_1 . Now assume that r is a slope of a side of the Newton polygon of

$A_{K_1}(L^d, M)$. The Newton polygon of $A_{K_1}(L^d, M)$ is obtained from the Newton polygon of $A_{K_1}(L, M)$ by simply replacing every vertex (i, j) with $(i/d, j)$. Consequently, dr is the slope of a side of the Newton polygon of $A_{K_1}(L, M)$, and therefore, is a strongly detected boundary slope of K_1 . $\square$

By work of Ohtsuki [15], all boundary slopes of 2-bridge knots are strongly detected. Thus, for 2-bridge knots we have a stronger result.

Corollary 2.5 *Suppose $K_1 \geq_d K_2$ and that K_1 is a 2-bridge knot. If $\{r_1, r_2, \dots, r_m\}$ is the set of boundary slopes of K_2 , then $\{dr_1, dr_2, \dots, dr_m\}$ is a subset of the set of boundary slopes of K_1 .*

For any knot K , the *diameter* of its set of boundary slopes is the maximum difference between any two slopes. We denote the diameter by $diam(K)$. For 2-bridge knots, in fact for all alternating Montesinos knots, $cr(K) = diam(K)/2$, where $cr(K)$ is the crossing number of K . See [9], [10] and [14]. This leads to the following results.

Corollary 2.6 *Suppose K_1 is a 2-bridge knot and $K_1 \geq_d K_2$. Then $cr(K_1) \geq |d|cr(K_2) \geq cr(K_2)$.*

Using Theorem 2.3(iii) and Corollary 2.6 it is now easy to show the following. (This result was first proven by Boileau, Rubinstein, and Wang [3].)

Corollary 2.7 *A 2-bridge knot can only be greater than or equal to finitely many other knots.*

3 2-bridge knots with four or less boundary slopes

In preparation for proving Theorem 4.2, we determine in this section those 2-bridge knots with small numbers of distinct boundary slopes. In [14], the authors determine necessary but not sufficient conditions on p and q so that $K_{p/q}$ has 4 or less distinct boundary slopes. In order to exploit Corollary 2.5 we require a stronger result than what appears in [14]. In particular, we need both a complete classification of 2-bridge knots with 4 or less boundary slopes as well as an explicit description of the associated slope sets.

We assume the reader is familiar with Hatcher and Thurston's paper [7] where a method for computing the boundary slopes of $K_{p/q}$ is given. We will also make use of a method




Figure 2: The portion of the Farey graph between $1/0$ and $7/17$.

equivalent to theirs which is described in [9]. Rather than describing the methodology in complete detail, we provide a brief description illustrated by an example.

Recall that the Farey graph may be thought of as the edges of the ideal modular tessellation of $\mathbb{H}^2$ obtained by starting with the ideal triangle whose vertices are $1/0$, $0/1$, and $1/1$ and then reflecting it in all possible ways across its edges. The ideal vertices of this tessellation are $\mathbb{Q} \cup \{\infty\}$. Two vertices a/b and c/d are joined by an edge precisely when $\det \begin{pmatrix} a & c \\ b & d \end{pmatrix} = \pm 1$ and we call this value the *determinant* of the directed edge from a/b to c/d . If a/b and c/d are joined by an edge, then so are a/b and $(a+c)/(b+d)$ as well as c/d and $(a+c)/(b+d)$. In this case the two fractions a/b and c/d , together with their *mediant* $(a+c)/(b+d)$, form the vertices of an ideal triangle in the tessellation. Finally, given any continued fraction expansion of p/q , the associated sequence of convergents defines a path in the Farey graph from $1/0$ to p/q . The number of triangles through which the path “turns” at each vertex is equal to the corresponding partial quotient.

For example, consider $7/17 = 0 + [2, 2, 3]$. We picture the relevant portion of the Farey graph in Figure 2. The sequence of convergents is $\{0, 1/2, 2/5, 7/17\}$. The partial quotients are 2, 2, 3 and the path turns through 2 triangles on the left at 0, 2 triangles on the right at $1/2$ and 3 triangles on the left at $2/5$. If all the partial quotients are positive, as in this example, then the turning at each vertex alternates between triangles on the left and right. Negative partial quotients correspond to turning in the opposite direction. Again, the reader is urged to consult [7] and [9].

For any p/q , there are infinitely many edge paths in the Farey graph that begin at $1/0$ and end at p/q , but only finitely many that are *minimal*. A path is minimal if it never backtracks and if it never traverses two edges of the same triangle. The minimal paths for $K_{7/17}$ and their corresponding continued fractions are shown in the first two columns of Table 1. In general, only one of the minimal paths is *even*, that is, all the

Table 1: Boundary slope data for $K_{7/17}$

minimal path γ	fraction	$m(\gamma)$	slope
$\{1/0, 1/1, 1/2, 3/7, 5/12, 7/17\}$	$1 + [-2, 4, -2, 2]$	4	0
$\{1/0, 0/1, 1/2, 3/7, 5/12, 7/17\}$	$0 + [2, 3, -2]$	2	4
$\{1/0, 1/1, 1/2, 2/5, 7/17\}$	$1 + [-2, 3, 3]$	1	6
$\{1/0, 0/1, 1/2, 2/5, 7/17\}$	$0 + [2, 2, 3]$	-1	10
$\{1/0, 1/3, 2/5, 7/17\}$	$0 + [3, -2, 4]$	-3	14

partial quotients are even. In Table 1, the first path is even. According to Hatcher and Thurston, each minimal path determines a boundary slope. The slope can be computed from the path γ by the formula

$$-2(m(\gamma) - m(\gamma_{\text{even}}))$$

where γ_{even} is the even path and $m(\gamma)$ is the sum of the determinants of each edge of the path, excluding the first edge. Thus, the even path always gives a slope of zero. For any path γ , we call $m(\gamma)$ the *unadjusted* slope. The last two columns of Table 1 give the unadjusted slopes and the boundary slopes of $K_{7/17}$.

For any p/q , if γ_1 and γ_2 are two paths from $1/0$ to p/q , then we can move from one path to the other by a sequence of *triangle moves*. By a triangle move, we mean replacing two consecutive edges that lie in a single triangle with the third edge of the triangle, or vice-versa. If γ_1 is changed to γ_2 by a single triangle move, and furthermore, the triangle lies on the right of γ_1 , then we call the move a *right triangle move*. Right triangle moves define a partial order on the set of paths: we say that $\gamma_1 > \gamma_2$ if γ_1 can be changed to γ_2 by a sequence of right triangle moves. Furthermore, it is shown in [9] that right triangle moves strictly decrease the unadjusted slope of the path, thus, $\gamma_1 > \gamma_2$ implies that $m(\gamma_1) > m(\gamma_2)$. The *upper* and *lower* minimal paths can now be defined as ones that are either maximal or minimal with respect to this partial order, respectively. These paths are well-defined because there is a unique minimal path with no triangles on its left and a unique minimal path with no triangles on its right. In Table 1, the first path is the upper path and the last path is the lower path.

The following lemma gives a lower bound on the number of distinct boundary slopes.

Lemma 3.1 *Suppose $p/q = r + [a_1, a_2, \dots, a_m]$ is a strongly positive continued fraction. Then the number of distinct boundary slopes of $K_{p/q}$ is at least $2 + \lfloor m/2 \rfloor$ and at most f_{m+1} where f_m denotes the m -th Fibonacci number (assuming $f_0 = 0$ and $f_1 = 1$).*


Proof. For the lower bound we will create a sequence of minimal paths $\gamma_{\text{upper}}, \gamma_1, \dots, \gamma_{\lfloor m/2 \rfloor}, \gamma_{\text{lower}}$ with strictly decreasing unadjusted slopes. Let γ_{upper} be the upper path. Next, let γ_1 be the path that goes from $1/0$ to $r/1$ to $r + [a_1]$ and then continues with the rest of the upper path. The path γ_1 is obtained from γ_{upper} by a sequence of a_1 right triangle moves. Now let γ_2 begin $\{1/0, r/1, r + [a_1, a_2], r + [a_1, a_2, a_2]\}$ and then continue with the rest of the upper path. Notice that γ_2 is obtained from γ_1 by a sequence of $a_2 + a_3$ right triangle moves. We continue in this manner to generate each path γ_i . The fact that both $a_1 > 1$ and $a_m > 1$ guarantee that all of these paths are minimal. Finally, their unadjusted slopes are strictly decreasing since each path is obtained from the previous one by performing right triangle moves. This establishes the lower bound. It is shown in [7] that f_m is an upper bound on the number of minimal paths, each of which might give a distinct boundary slope. $\square$

As the number m of partial quotients grows, it becomes tedious to characterize all knots with m partial quotients and exactly k distinct boundary slopes. Therefore, in the following theorem, we only classify 2-bridge knots with up to four distinct boundary slopes. Furthermore, we provide the set of slopes for only one knot from each chiral pair. (Remember that switching from a knot to its mirror image will negate the set of boundary slopes.)

Theorem 3.2 *Let $K_{p/q}$ be a 2-bridge knot with 2, 3, or 4, distinct boundary slopes. Then a strongly positive continued fraction representing $K_{p/q}$, or its mirror image, together with the associated slope set, is given in the following table.*

	continued fraction	slope set *=multiplicity 2, **=multiplicity 3
2	$[a_1], a_1 \text{ odd}$	$\{0, 2a_1\}$
3_i	$[a_1, a_2], a_1 \text{ even}, a_2 \text{ even}$	$\{-2a_1, 0, 2a_2\}$
3_{ii}	$[a_1, a_2], a_1 \text{ even}, a_2 \text{ odd}$	$\{0, 2a_1, 2a_1 + 2a_2\}$
3_{iii}	$[a_1, 1, a_1], a_1 \text{ odd}$	$\{-4a_1 - 2, -2a_1 - 2^*, 0\}$
4_i	$[a_1, a_2, a_1], a_1 \text{ odd}, a_2 \text{ odd}, a_2 > 1$	$\{-4a_1 - 2a_2, -2a_1 - 2a_2^*, -2a_2, 0\}$
4_{ii}	$[a_1, 1, a_3], a_1 \neq a_3, a_1 \text{ odd}, a_3 \text{ odd}$	$\{-2a_1 - 2a_3 - 2, -2a_3 - 2, -2a_1 - 2, 0\}$
4_{iii}	$[a_1, 1, a_3], a_1 \neq a_3, a_1 \text{ even}, a_3 \text{ odd}$	$\{-2a_1, 0, -2a_1 + 2a_3, 2a_3 + 2\}$
4_{iv}	$[a_1, 1, a_1, a_1 + 1], a_1 \text{ even}$	$\{-2a_1, 0^*, 2a_1 + 2^*, 4a_1 + 4\}$
4_v	$[2, 1, 1, 1, 2] = 8/21$	$\{-8, -4^*, 0^{**}, 6\}$

Proof. Suppose that $K_{p/q}$ has 4 or less distinct boundary slopes and strongly positive continued fraction expansion $p/q = 0 + [a_1, a_2, \dots, a_m]$. By Lemma 3.1, it follows that

Figure 3: The partially ordered set of minimal paths when $m = 4$.

$m \leq 4$. We will examine only the case $m = 3$ and leave the remaining cases, which are handled in a similar manner, to the interested reader.

Figure 3 shows all possible minimal paths for $p/q = [a_1, a_2, a_3]$ arranged according to the partial ordering on paths. Each of the five figures is made up of three main triangles which are subdivided into a_1 , a_2 , and a_3 smaller triangles, respectively, as we move from left to right. The five paths γ_{upper} , γ_1 , γ_2 , γ_3 , and γ_{lower} are depicted with heavy dark lines. Also listed is the unadjusted slope m for each path under the path's label.

Since $a_1 > 1$ and $a_3 > 1$ we have that γ_{upper} , γ_1 , γ_2 , and γ_{lower} are minimal. If γ_3 is minimal and $a_1 \neq a_3$, then we would have 5 distinct boundary slopes. Therefore, we have two cases to consider.

Assume γ_3 is not minimal, which requires $a_2 = 1$, and also that $a_1 \neq a_3$. Thus, $p/q = 0 + [a_1, 1, a_3]$ and we have four distinct boundary slopes. In order to compute

the (adjusted) slopes we need to determine the unique even path. Notice that

$$\frac{p}{q} = [a_1, 1, a_3] = \frac{1 + a_3}{a_1 + a_3 + a_1 a_3}$$

and so in order for q to be odd we need at least one of a_1 or a_3 to be odd. This gives three sub-cases which are listed below.

a_1	a_3	even path	boundary slopes
even	odd	γ_1	$\{-2a_1, 0, -2a_1 + 2a_3, 2a_3 + 2\}$
odd	even	γ_2	$\{-2a_3, 2a_1 - 2a_3, 0, 2a_1 + 2\}$
odd	odd	γ_{lower}	$\{-2a_1 - 2a_3 - 2, -2a_3 - 2, -2a_1 - 2, 0\}$

These slopes appear in parts 4_{ii} and 4_{iii} of Theorem 3.2. The first two rows correspond to equivalent knots and only appear in one row of the theorem.

Next assume that γ_3 is not minimal and $a_1 = a_3$. In this case, we have 3 distinct boundary slopes and

$$\frac{p}{q} = [a_1, 1, a_1] = \frac{1 + a_1}{a_1^2 + 2a_1}.$$

Here q is odd if and only if a_1 is odd. In this case, the even path is γ_{lower} and the boundary slopes are

$$\{-4a_1 - 2, -2a_1 - 2, 0\}$$

where the second slope $-2a_1 - 2$ has multiplicity two, that is, it corresponds to two paths, γ_1 and γ_2 . This data appears in part 3_{iii} of Theorem 3.2.

Finally, assume that γ_3 is minimal, which means $a_2 > 1$, and $a_1 = a_3$. We now have four distinct boundary slopes and

$$\frac{p}{q} = [a_1, a_2, a_1] = \frac{1 + a_1 a_2}{a_1^2 a_2 + 2a_1}.$$

In order for $q = a_1^2 a_2 + 2a_1$ to be odd we must have that both a_1 and a_2 are odd. This implies that γ_{lower} is the even path and we obtain the following boundary slopes:

$$\{-4a_1 - 2a_2, -2a_1 - 2a_2, -2a_2, 0\}.$$

Notice that the second boundary slope $-2a_1 - 2a_2$ has multiplicity two. This data appears in part 4_i of Theorem 3.2. The remaining cases follow by a similar analysis with $m = 1$, $m = 2$, or $m = 4$. $\square$

By working out the fractions p/q for each case of Theorem 3.2 we obtain the following corollary which strengthens Theorem 3 of [14]. As in Theorem 3.2, we consider a knot and its mirror image to be equivalent.

Corollary 3.3 *Let K be a 2-bridge knot.*

- i. *K has exactly two distinct boundary slopes if and only if $K \equiv K_{1/q}$.*
- ii. *K has exactly three distinct boundary slopes if and only if K is not equivalent to a knot from part (i) and $K \equiv K_{p/q}$ for which either*
 - a. *$p \mid q - 1$ or*
 - b. *$p^2 = q + 1$.*
- iii. *K has exactly four distinct boundary slopes if and only if K is not equivalent to a knot from parts (i) or (ii) and $K \equiv K_{p/q}$ for which either*
 - a. *$p + 1 \mid q$ and $q \mid p^2 - 1$,*
 - b. *$p \mid q + 1$,*
 - c. *$(p - 1)^3 = q^2$, or*
 - d. *$p/q = 8/21$.*

4 Ohtsuki, Riley, Sakuma knot pairs

In this section we briefly review the method of Ohtsuki, Riley, and Sakuma for constructing a pair of two bridge knots $K_{p'/q'}$ and $K_{p/q}$ with $K_{p'/q'} \geq K_{p/q}$. The reader is encouraged to consult [16] for a more detailed description. One begins with any 2-bridge knot (or link) $K_{p/q}$ given by the four-plat shown in Figure 4. The 4-string braid defining $K_{p/q}$ is denoted by β . There are three associated braids β^{-1} , β_- , and β_-^{-1} which are obtained by rotating β 180 degrees around an axis perpendicular to the plane of the diagram, reflecting β through a plane perpendicular to the plane of the diagram, and the composition of these two motions, respectively. In order to construct a 2-bridge knot $K_{p'/q'}$ that is greater than $K_{p/q}$ we consider a 4-plat with an odd number of “boxes” each containing either β , β^{-1} , β_- , or β_-^{-1} . Starting from the left in Figure 4, the first box contains β and every other box after that contains β or β_- . The remaining boxes each contain β^{-1} or β_-^{-1} . Also between each braid box in $K_{p'/q'}$ we may insert an even number of half-twists in the middle two strands of the 4-plat. Finally, a *branched fold map* f is constructed from the complement of $K_{p'/q'}$ onto the complement of $K_{p/q}$ as follows. The complement of $K_{p'/q'}$ is cut along a collection of parallel 2-spheres into two 3-balls and a number of $S^2 \times I$ ’s. The complement of $K_{p/q}$ is decomposed into two 3-balls and a single $S^2 \times I$. A continuous mapping is then defined in a piecewise manner. First each 3-ball “upstairs” is mapped by the identity onto a corresponding 3-ball “downstairs.” Next, each $S^2 \times I$ containing one braid box




Figure 4: The Ohtsuki, Riley, Sakuma construction.

upstairs is mapped homeomorphically onto the $S^2 \times I$ downstairs in a way that depends on the presence of β , β^{-1} , β_- , or β_-^{-1} . Finally, the mapping is extended to the remaining $S^2 \times I$'s upstairs onto the 3-balls downstairs using 2-fold branched mappings that depend on the combinations of β , β^{-1} , β_- , or β_-^{-1} in the adjacent components. From the way that f is constructed, the two meridional generators associated with the bridges at either end of the 4-plat diagram upstairs are taken to the corresponding generators downstairs. Furthermore, the longitude upstairs is taken to a power of the longitude downstairs. Thus f induces an epimorphism on fundamental groups which preserves peripheral structure.

Given a 2-bridge knot $K_{p/q}$, Ohtsuki, Riley and Sakuma determine which knots $K_{p'/q'}$ are produced by their construction. In particular, they show that p'/q' gives such a knot if and only if it lies in the orbit of p/q under the action of $\Gamma_\infty * \Gamma_{p/q}$, where $\Gamma_{r/s}$ is the infinite dihedral group generated by all reflections in edges of the Farey graph which end at r/s . Let A_1 and A_2 be generators of Γ_∞ defined by reflection in the edges $\langle 1/0, 0/1 \rangle$ and $\langle 1/0, 1/1 \rangle$, respectively. For $p/q = [a_1, a_2, \dots, a_m]$ let B_1 and B_2 be generators of $\Gamma_{p/q}$ defined by reflection in the edges $\langle p/q, [a_1, a_2, \dots, a_{m-1}] \rangle$ and $B_2 = \langle p/q, [a_1, a_2, \dots, a_m - 1] \rangle$. Thus $p'/q' = W(p/q)$ where W is a word in A_1, A_2, B_1 and B_2 . Since B_1 and B_2 fix p/q , and A_1 and A_2 take p/q to a fraction representing the same knot, we may assume that we begin by applying a nontrivial word in A_1 and A_2 and end by applying a nontrivial word in B_1 and B_2 . Hence we

may write W as

$$(3) \quad W = W_1 W_2 \dots W_n, \text{ with } n \text{ even,}$$

where

$$(4) \quad W_i = \begin{cases} (B_1 B_2)^{c_i} B_1^{\frac{1-\eta_i}{2}}, & c_i \in \mathbb{Z}, \eta_i \in \{1, -1\}, \text{ if } i \text{ is odd, and} \\ (A_1 A_2)^{c_i} A_1^{\frac{1-\eta_i}{2}}, & c_i \in \mathbb{Z}, \eta_i \in \{1, -1\}, \text{ if } i \text{ is even.} \end{cases}$$

Ohtsuki, Riley and Sakuma then show (Lemma 5.3 in [16]) that

$$(5) \quad p'/q' = [\epsilon_1 \mathbf{a}, 2\epsilon_1 c_1, \epsilon_2 \mathbf{a}^{-1}, 2\epsilon_2 c_2, \epsilon_3 \mathbf{a}, 2\epsilon_3 c_3, \epsilon_4 \mathbf{a}^{-1}, \dots, \epsilon_{n+1} \mathbf{a}],$$

where $\mathbf{a}$ is the vector $\mathbf{a} = (a_1, a_2, \dots, a_m)$, $\mathbf{a}^{-1}$ means $\mathbf{a}$ written backwards, $\epsilon_1 = 1$, and $\epsilon_i = \prod_{j=1}^{i-1} -\eta_j$. The following lemma will prove helpful in simplifying continued fractions.

Lemma 4.1 *Let $\mathbf{a}$ and $\mathbf{b}$ be vectors of integers, possibly empty, and m and n be any integers. Then*

- i. $r + [\mathbf{a}, m, -n, \mathbf{b}] = r + [\mathbf{a}, m-1, 1, n-1, -\mathbf{b}]$ and
- ii. $r + [\mathbf{a}, m, 0, n, \mathbf{b}] = r + [\mathbf{a}, m+n, \mathbf{b}]$

Proof. If $p/q = r_0 + [r_1, r_2, \dots, r_k]$ then it is well known that

$$\pm \begin{pmatrix} p \\ q \end{pmatrix} = M_{(r_0, r_1, \dots, r_k)} \begin{pmatrix} 1 \\ 0 \end{pmatrix},$$

where

$$M_{(r_0, r_1, \dots, r_k)} = \begin{pmatrix} r_0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} r_1 & 1 \\ 1 & 0 \end{pmatrix} \cdots \begin{pmatrix} r_k & 1 \\ 1 & 0 \end{pmatrix}.$$

It is an easy exercise to show that

$$\begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} M_{\mathbf{b}} = (-I)^{|\mathbf{b}|} M_{-\mathbf{b}} \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix},$$

where $|\mathbf{b}|$ is the dimension of the vector $\mathbf{b}$ and I is the identity matrix. The proof of Lemma 4.1(i) follows from

$$\begin{aligned}
M_{\mathbf{a},m,-n,\mathbf{b}} \begin{pmatrix} 1 \\ 0 \end{pmatrix} &= M_{\mathbf{a}} \begin{pmatrix} m & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} -n & 1 \\ 1 & 0 \end{pmatrix} M_{\mathbf{b}} \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\
&= M_{\mathbf{a}} \begin{pmatrix} m-1 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} n-1 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} M_{\mathbf{b}} \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\
&= (-I)^{|\mathbf{b}|} M_{\mathbf{a},m-1,1,n-1,-\mathbf{b}} \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\
&= (-I)^{|\mathbf{b}|} M_{\mathbf{a},m-1,1,n-1,-\mathbf{b}} \begin{pmatrix} -1 \\ 0 \end{pmatrix} \\
&= (-I)^{|\mathbf{b}|+1} M_{\mathbf{a},m-1,1,n-1,-\mathbf{b}} \begin{pmatrix} 1 \\ 0 \end{pmatrix}
\end{aligned}$$

The second part of the lemma follows in a similar way. $\square$

We are now prepared to show that in any ORS pair $(K_{p'/q'}, K_{p/q})$, if $K_{p'/q'}$ is not a torus knot, then it has at least five distinct boundary slopes.

Theorem 4.2 *If $K_{p'/q'} \geq K_{p/q}$ is a nontrivial ORS pair, then either*

- i. $K_{p'/q'}$ and $K_{p/q}$ are both torus knots and $K_{p'/q'}$ has precisely two distinct boundary slopes, or
- ii. $K_{p'/q'}$ has at least five distinct boundary slopes.

Proof. Let $p/q = [a_1, a_2, \dots, a_m] = [\mathbf{a}]$ be a strongly positive continued fraction. Suppose that $p'/q' = W(p/q)$ where W is described in Equations (3) and (4). By Equation (5) we have that

$$(6) \quad p'/q' = [\epsilon_1 \mathbf{a}, 2\epsilon_1 c_1, \epsilon_2 \mathbf{a}^{-1}, 2\epsilon_2 c_2, \epsilon_3 \mathbf{a}, 2\epsilon_3 c_3, \epsilon_4 \mathbf{a}^{-1}, \dots, \epsilon_{n+1} \mathbf{a}].$$

Suppose $c_i = 0$ for some i . Because W_i is not the identity, it follows that $\eta_i = -1$ and hence $\epsilon_i = \epsilon_{i+1}$. Therefore, on either side of $2\epsilon_i c_i = 0$, the continued fraction appears as $p'/q' = [\dots, \epsilon_i \mathbf{a}^{\pm 1}, 0, \epsilon_i \mathbf{a}^{\mp 1}, \dots]$. Using Lemma 4.1, we may eliminate the zero to combine $\epsilon_i \mathbf{a}^{\pm 1}$ and $\epsilon_i \mathbf{a}^{\mp 1}$ into a single strongly positive or negative vector.

Let $1 \leq j_1 < j_2 < \dots < j_k \leq n$ be the indices of the nonzero c_i 's. Eliminating each of the $n - k$ zero entries as just described gives the following continued fraction expansion with $n(m-1) + m + 2k$ partial quotients,

$$(7) \quad p'/q' = [\epsilon_1 \mathbf{v}_1, 2\epsilon_{j_1} c_{j_1}, \epsilon_{j_2} \mathbf{v}_2, 2\epsilon_{j_2} c_{j_2}, \epsilon_{j_3} \mathbf{v}_3, \dots, 2\epsilon_{j_k} c_{j_k}, \epsilon_{n+1} \mathbf{v}_{k+1}]$$

where $\epsilon_1 = 1$, each $\mathbf{v}_i$ is a strongly positive vector, and $c_{j_i} \neq 0$ for all i . Notice that if $m = 1$, then $\mathbf{a}^{-1} = \mathbf{a} = a_1$ and each vector $\mathbf{v}_i$ is a single integer given by $\mathbf{v}_i = (j_i - j_{i-1})a_1$ if $i \leq k$ and $\mathbf{v}_{k+1} = (n + 1 - j_k)a_1$. Otherwise, if $m > 1$, $|\mathbf{v}_i| \geq 2$ for all i .

We now claim that the continued fraction given in Equation (7) can be changed into a strongly positive one with at least as many partial quotients. Before proving this claim, consider its consequences. If $m = 1$ then the number of partial quotients is at least $2k + 1$ which can only be less than 6 if $k = 0, 1$, or 2 . If $m > 1$, the number of partial quotients can only be less than 6 if $n = m = 2$ and $k = 0$. Hence, except in the cases where $m = 1$ and $0 \leq k \leq 2$, or $n = m = 2$ and $k = 0$, we can express p'/q' as a strongly positive continued fractions with at least 6 partial quotients. Lemma 3.1 now implies that $K_{p'/q'}$ has at least 5 distinct boundary slopes. Thus, after proving this claim, we must analyze these four cases.

To prove the claim, we first assume that $m > 1$ and proceed by induction on the number of sign changes in the sequence of partial quotients in Equation (7). If there are none, then it is easy to check that Equation (7) is already strongly positive. If there are sign changes, then consider the smallest value of r so that $\epsilon_{j_r} = 1$ and either c_{j_r} or $\epsilon_{j_{r+1}}$ is negative.

If $c_{j_r} > 0$ and $\epsilon_{j_{r+1}} = -1$, then using Lemma 4.1, we obtain

$$\begin{aligned} p'/q' &= [\dots, \mathbf{v}_r, 2c_{j_r}, -\mathbf{v}_{r+1}, -2c_{j_{r+1}}, \dots] \\ &= [\dots, \mathbf{v}_r, 2c_{j_r}, -v_{r+1}^1, -v_{r+1}^2, \dots, -v_{r+1}^{|\mathbf{v}_{r+1}|}, -2c_{j_{r+1}}, \dots] \\ &= [\dots, \mathbf{v}_r, 2c_{j_r} - 1, 1, v_{r+1}^1 - 1, v_{r+1}^2, \dots, v_{r+1}^{|\mathbf{v}_{r+1}|}, 2c_{j_{r+1}}, \dots]. \end{aligned}$$

This continued fraction is now of the same form as the original, has fewer sign changes, and has more partial quotients. The result now follows by induction. It is important here to remember that since $m > 1$, $|\mathbf{v}_i| \geq 2$ for all i .

If $c_{j_r} < 0$ and $\epsilon_{j_{r+1}} = -1$, then using Lemma 4.1, we obtain

$$\begin{aligned} p'/q' &= [\dots, \mathbf{v}_r, 2c_{j_r}, -\mathbf{v}_{r+1}, -2c_{j_{r+1}}, \dots] \\ &= [\dots, v_r^1, v_r^2, \dots, v_r^{|\mathbf{v}_r|}, 2c_{j_r}, -\mathbf{v}_{r+1}, -2c_{j_{r+1}}, \dots] \\ &= [\dots, v_r^1, v_r^2, \dots, v_r^{|\mathbf{v}_r|} - 1, 1, -2c_{j_r} - 1, \mathbf{v}_{r+1}, 2c_{j_{r+1}}, \dots]. \end{aligned}$$

Again, this continued fraction is now of the same form as the original, has fewer sign changes, and has more partial quotients. The result now follows by induction.

Finally, suppose that $c_{j_r} < 0$ and $\epsilon_{j_{r+1}} = 1$. Applying Lemma 4.1 twice, we obtain

$$\begin{aligned} p'/q' &= [\dots, \mathbf{v}_r, 2c_{j_r}, \mathbf{v}_{r+1}, \dots] \\ &= [\dots, v_r^1, \dots, v_r^{|\mathbf{v}_r|} - 1, 1, -2c_{j_r} - 2, 1, v_{r+1}^1 - 1, v_{r+1}^2, \dots, v_{r+1}^{|\mathbf{v}_{r+1}|}, \dots]. \end{aligned}$$

If $c_{j_r} \neq -1$, then once again we have arrived a continued fraction of the same form as the original, but with fewer sign changes and more partial quotients. If instead, $c_{j_r} = -1$, then we may use Lemma 4.1 to eliminate the zero entry and arrive at a continued fraction with the same form as the original, fewer sign changes, but now an equal number of partial quotients. In either case, the result follows by induction.

We now consider the case where $m = 1$. Recall that if $m = 1$, then $\mathbf{a}^{-1} = \mathbf{a} = a_1$, $\mathbf{v}_i = (j_i - j_{i-1})a_1$ for $i \leq k$, and $\mathbf{v}_{k+1} = (n + 1 - j_k)a_1$. Equation (7) is now of the form

$$(8) \quad p'/q' = [\epsilon_1 j_1 a_1, 2\epsilon_{j_1} c_{j_1}, \epsilon_{j_2} (j_2 - j_1) a_1, 2\epsilon_{j_2} c_{j_2}, \dots, 2\epsilon_{j_k} c_{j_k}, \epsilon_{n+1} (n + 1 - j_k) a_1].$$

Because $a_1 \geq 3$, this continued fraction is a member of a family $\mathcal{F}$ of continued fractions all having the form $[\mathbf{v}, \mathbf{w}]$ where

- $\mathbf{v}$ is a strongly positive vector (possibly of length 1),
- the first entry of the vector $\mathbf{w}$ is negative,
- the entries of $\mathbf{w}$ alternate between being even and having magnitudes greater than 2 (the first entry of $\mathbf{w}$ can have either property), and
- the last entry of $\mathbf{w}$ has magnitude greater than 2.

We call $\mathbf{v}$ the *strongly positive part* and $\mathbf{w}$ the *tail* of such a continued fraction. We shall prove, by induction on the length of the tail, that any such continued fraction can be changed to a strongly positive one with at least as many partial quotients. Since $\mathbf{v}$ is strongly positive, if $\mathbf{w}$ has length zero then we are done. Now consider such a continued fraction where $\mathbf{w}$ is not empty. We have

$$\begin{aligned} [\mathbf{v}, \mathbf{w}] &= [v_1, v_2, \dots, v_r, w_1, w_2, \dots, w_s] \\ &= [v_1, v_2, \dots, v_r - 1, 1, -w_1 - 1, -w_2, \dots, -w_s] \end{aligned}$$

If $-w_2 > 0$, or if $-w_2 < 0$ and $-w_1 - 1 > 1$, then we have arrived at a member of $\mathcal{F}$ with more partial quotients and a shorter tail. Hence the result follows by induction. Otherwise, we have $-w_1 - 1 = 1$ and $-w_2 < 0$. Thus

$$\begin{aligned} [\mathbf{v}, \mathbf{w}] &= [v_1, v_2, \dots, v_r, -2, w_2, \dots, w_s] \\ &= [v_1, v_2, \dots, v_r - 1, 1, 1, -w_2, \dots, -w_s] \\ &= [v_1, v_2, \dots, v_r - 1, 1, 0, 1, w_2 - 1, w_3, \dots, w_s] \\ &= [v_1, v_2, \dots, v_r - 1, 2, w_2 - 1, w_3, \dots, w_s]. \end{aligned}$$

Because the entries of $\mathbf{w}$ alternate between being even and having magnitudes greater than 2, it must be the case that $w_2 > 2$ and hence $w_2 - 1 > 1$. Hence we have arrived

at a member of $\mathcal{F}$ with the same number of partial quotients but with a shorter tail. Once again, the result now follows by induction.

We have now completed the proof of the claim, and it remains to analyze the four cases already mentioned above. If $m = n = 2$ and $k = 0$, we have $p/q = [a_1, a_2]$ and $p'/q' = [a_1, a_2, 0, a_2, a_1, 0, a_1, a_2] = [a_1, 2a_2, 2a_1, a_2]$. Comparing this strongly positive continued fraction to the table in Theorem 3.2 we see that $K_{p'/q'}$ cannot have less than 5 distinct boundary slopes.

If $m = 1$ and $k = 0$ the continued fraction for p'/q' collapses to $p'/q' = [(n+1)a_1]$. Thus $K_{p'/q'}$ is a torus knot with exactly 2 slopes.

Next, suppose that $m = 1$ and $k = 1$. Then exactly one of the c_i 's is nonzero, say c_{j_1} . Equation 7 now gives $p'/q' = [j_1a_1, 2\epsilon_{j_1}c_{j_1}, \epsilon_{n+1}(n+1-j_1)a_1]$. If all of these partial quotients are positive, then comparing this strongly positive continued fraction to the table in Theorem 3.2, we see that $K_{p'/q'}$ cannot have less than 5 distinct boundary slopes. If not, there are either one or two sign changes in the sequence of partial quotients. These considerations lead to the following cases which are handled in a fashion similar to that when $m > 1$:

- If $\epsilon_{j_1}c_{j_1} > 0$ and $\epsilon_{n+1} = -1$, then $p'/q' = [j_1a_1, 2\epsilon_{j_1}c_{j_1} - 1, 1, (n+1-j_1)a_1 - 1]$.
- If $\epsilon_{j_1}c_{j_1} < 0$ and $\epsilon_{n+1} = -1$, then $p'/q' = [j_1a_1 - 1, 1, -2\epsilon_{j_1}c_{j_1} - 1, (n+1-j_1)a_1]$.
- If $\epsilon_{j_1}c_{j_1} = -1$ and $\epsilon_{n+1} = 1$, then $p'/q' = [j_1a_1 - 1, 2, (n+1-j_1)a_1 - 1]$.
- If $\epsilon_{j_1}c_{j_1} < -1$ and $\epsilon_{n+1} = 1$, then $p'/q' = [j_1a_1 - 1, 1, -2\epsilon_{j_1}c_{j_1} - 2, 1, (n+1-j_1)a_1 - 1]$.

In all four of these cases, comparison to the table in Theorem 3.2 gives that $K_{p'/q'}$ cannot have less than 5 distinct boundary slopes.

Our final case to consider is when $m = 1$ and $k = 2$. Now two of the c_i 's are nonzero, say c_{j_1} and c_{j_2} . Eliminating zeroes leads to

$$p'/q' = [j_1a_1, 2\epsilon_{j_1}c_{j_1}, \epsilon_{j_2}(j_2 - j_1)a_1, 2\epsilon_{j_2}c_{j_2}, \epsilon_{n+1}(n+1-j_2)a_1].$$

If all of these partial quotients are positive, then comparing this strongly positive continued fraction to the table in Theorem 3.2, we once again see that $K_{p'/q'}$ cannot have less than 5 distinct boundary slopes. If not, then using Lemma 4.1 to move to a strongly positive expansion will increase the number of partial quotients to 6 or more, again resulting in 5 or more boundary slopes, unless the first entry of the tail is -2 . In this case, that entry will remain even, and comparison with the table in Theorem 3.2 again shows that this knot cannot have less than 5 distinct boundary slopes. $\square$

The lower bound of five boundary slopes in part *ii* of Theorem 4.2 is sharp. For example, with $p/q = 1/3 = [3]$ and $p'/q' = 7/45 = [3, 0, 3, 2, 3] = [6, 2, 3]$ we have $K_{7/45} \geq K_{1/3}$. Furthermore, using Figure 3, it is easy to show that $K_{7/45}$ has exactly 5 distinct boundary slopes.

Recall that if $K_{p'/q'}$ is a 2-bridge knot and $K_{p'/q'} \geq K_2$, then K_2 is also a 2-bridge knot. Combining this fact with the result in Theorem 4.2 gives the following corollary.

Corollary 4.3 *If the answer to Question 1.1 is yes, then 2-bridge knots with exactly 3 or 4 distinct boundary slopes are minimal with respect to the Silver-Whitten partial ordering.*

Notice that if K is a 2-bridge knot with exactly 2 distinct boundary slopes, then K is the torus knot $T_{2,q}$ and is minimal if and only if q is prime.

5 Two-bridge knots with three distinct boundary slopes

In this section we provide evidence for an affirmative answer to Question 1.1 by proving that 2-bridge knots with exactly three distinct boundary slopes are in fact minimal.

Theorem 5.1 *If K is a 2-bridge knot with exactly 3 distinct boundary slopes, then K is minimal with respect to the Silver-Whitten partial order.*

Proof. Suppose K_1 is a 2-bridge knot with exactly three distinct boundary slopes and that $K_1 \geq_d K_2$ with K_2 nontrivial. Then, by Theorem 2.3 and Corollary 2.5, K_2 is a 2-bridge knot with exactly two or three distinct boundary slopes. If K_2 has two distinct slopes then it is a torus knot and so, by González-Acuña and Ramírez [6], K_1 is given by the ORS construction. Now Theorem 4.2 contradicts that K_1 has exactly three boundary slopes. Therefore, from Theorem 3.2 there are three possibilities for K_1 and three for K_2 , giving a total of 9 different cases to consider. In what follows, the symbol $(3_{ii}, 3_i)$, for example, will be used to denote the case where K_1 is the second type of knot with three slopes and K_2 is the third type of knot with three slopes, as listed in Theorem 3.2.

Since boundary slopes of 2-bridge knots are always even, we first consider them mod 4. If K_2 has r distinct slopes equal to 0 mod 4 and s distinct slopes equal to 2 mod 4, then K_1 has the same number of slopes of each type because of Corollary 2.5 and the fact that d is odd. This observation rules out four cases: $(3_i, 3_{ii})$, $(3_i, 3_{iii})$, $(3_{ii}, 3_i)$, $(3_{iii}, 3_i)$.

Before considering any other cases we note that the Alexander polynomial of a 2-bridge knot with three boundary slopes is given in the following table. Note that in each case, no nontrivial constant can be factored from each polynomial.

knot type	Alexander polynomial
3_i $0 + [a_1, a_2]$, a_1 even, a_2 even	$-\frac{a_1 a_2}{4} + \left(1 - \frac{a_1 a_2}{2}\right)t - \frac{a_1 a_2}{4}t^2$
3_{ii} $0 + [a_1, a_2]$, a_1 even, a_2 odd	$\frac{a_2 + 1}{2} + a_2 \sum_{i=1}^{a_1-1} (-1)^i t^i + \frac{a_2 + 1}{2} t^{a_1}$
3_{iii} $0 + [a_1, -1, a_1]$, a_1 odd	$\frac{(a_1 + 1)^2}{4} + \left(1 - \frac{(a_1 + 1)^2}{2}\right)t + \frac{(a_1 + 1)^2}{4}t^2$

Case $(3_i, 3_i)$ Suppose K_1 corresponds to $0 + [a_1, a_2]$ and K_2 corresponds to $0 + [b_1, b_2]$. Because of Corollary 2.5, it follows that

$$\{-2db_1, 0, 2db_2\} = \{-2a_1, 0, 2a_2\}$$

and so $a_1 a_2 = d^2 b_1 b_2$. Furthermore, Δ_{K_2} divides Δ_{K_1} which implies that $d = \pm 1$. Hence, $\{a_1, a_2\} = \{b_1, b_2\}$ and we have $K_2 \equiv K_1$.

Case $(3_{ii}, 3_{iii})$ Suppose K_1 corresponds to $0 + [a_1, a_2]$ and K_2 corresponds to $0 + [b_1, 1, b_1]$. From Corollary 2.5 we obtain that $d < 0$, $a_1 = -(b_1 + 1)d$, and $a_1 + a_2 = -(2b_1 + 1)d$. From these it follows that $a_2 = -b_1 d$. Since the determinants of these knots are $a_1 a_2 + 1$ and $b_1^2 + 2b_1$, respectively, it follows from Theorem 2.3 that $k(b_1^2 + 2b_1) = a_1 a_2 + 1 = b_1^2 d^2 + b_1 d^2 + 1$ for some integer k . Hence b_1 divides 1 which is a contradiction.

Case $(3_{iii}, 3_{ii})$ Suppose K_1 corresponds to $0 + [a_1, 1, a_1]$ and K_2 corresponds to $0 + [b_1, b_2]$. From Corollary 2.5 we obtain that $d < 0$, $a_1 + 1 = -b_1 d$, and $2a_1 + 1 = -(b_1 + b_2)d$. From these we obtain that $d = -1$, $b_1 = a_1 + 1$, and $b_2 = a_1$. However, the division of the Alexander polynomials implies that $b_1 = 2$ and this contradicts that $a_1 > 1$.

Case $(3_{iii}, 3_{iii})$ Suppose K_1 corresponds to $0 + [a_1, 1, a_1]$ and K_2 corresponds to $0 + [b_1, 1, b_2]$. The division of the Alexander polynomials implies that $b_1 = a_1$ and hence $K_2 = K_1$.

Case (3_{ii}, 3_{ii}) Suppose K_1 corresponds to $0 + [a_1, a_2]$ and K_2 corresponds to $0 + [b_1, b_2]$. Using Corollary 2.5 we immediately obtain that $a_1 = b_1d$ and $a_2 = b_2d$. All evidence suggests that if $d \neq 1$, then the Alexander polynomials do not divide (in fact, they are almost certainly both irreducible), however, we were unable to prove this in general. Instead we turn to character varieties to settle this last case.

By Theorem 2.3, we have that $\phi^* : X(K_2) \rightarrow X(K_1)$ is an injective, algebraic and closed (in the Zariski topology) mapping. In particular, this implies that the image $\phi^*(X(K_2))$ is a subvariety of $X(K_1)$ that is birationally equivalent to $X(K_2)$. By Theorem 6.5 of Macasieb, Petersen, and Van Luijk [13], both $X(K_1)$ and $X(K_2)$ are irreducible curves and so we conclude that $X(K_1)$ and $X(K_2)$ are birationally equivalent, and therefore have the same genus. Moreover, by Theorem 6.5 of [13], the genus of $X(K_2)$ is

$$(9) \quad 3 \left(\frac{b_2 + 1}{2} \right) \left(\frac{b_1}{2} \right) - \frac{b_2 + 1}{2} - 4 \frac{b_1}{2} + 2.$$

(To interpret Theorem 6.5 in our setting, note that $k = -b_2$ and $l = b_1$.) Similarly, the genus of $X(K_1)$ is

$$(10) \quad 3 \left(\frac{db_2 + 1}{2} \right) \left(\frac{db_1}{2} \right) - \frac{db_2 + 1}{2} - 4 \frac{db_1}{2} + 2.$$

Equating (9) and (10) gives

$$d(3db_1b_2 - 5b_1 - 2b_2) = 3b_1b_2 - 5b_1 - 2b_2.$$

Since d , b_1 , and b_2 are all positive, this can only be true if $d = 1$, in which case $K_2 = K_1$.

□

Since Theorem 3.2 also classifies those 2-bridge knots with exactly four distinct boundary slopes, we could hope to apply arguments similar to those above to prove that 2-bridge knots with four slopes are minimal. In particular, looking at boundary slopes will eliminate most of the 40 cases involved. However, without additional information about the relevant Alexander polynomials or character varieties, we were unable to settle resolve several cases. We close with the following conjecture and question.

Conjecture 5.2 *A 2-bridge knot with exactly four distinct boundary slopes is minimal with respect to the Silver-Whitten partial ordering.*

Question 5.3 *Does there exist a non 2-bridge, non-minimal knot with exactly 3 distinct boundary slopes?*

References

- [1] **M Boileau, S Boyer**, *On character varieties, sets of discrete characters, and non-zero degree maps* (2007) ArXiv:math.GT/0701384
- [2] **M Boileau, S Boyer, R A, W S**, *Simon's conjecture for 2-bridge knots* (2009) ArXiv:math.GT/09032898
- [3] **M Boileau, J H Rubinstein, S Wang**, *Finiteness of 3-manifolds associated with non-zero degree mappings* (2005) ArXiv:math/0511541
- [4] **G Burde, H Zieschang**, *Knots*, volume 5 of *de Gruyter Studies in Mathematics*, Walter de Gruyter & Co., Berlin (2003)
- [5] **D Cooper, M Culler, H Gillet, D D Long, P B Shalen**, *Plane curves associated to character varieties of 3-manifolds*, Invent. Math. 118 (1994) 47–84
- [6] **F González-Acuña, A Ramírez**, *Two-bridge knots with property Q*, Quart. J. Math. 52 (2001) 447–454
- [7] **A Hatcher, W Thurston**, *Incompressible surfaces in 2-bridge knot complements*, Invent. Math. 79 (1985) 225–246
- [8] **J Hoste, P D Shanahan**, *A formula for the A-polynomial of twist knots*, J. Knot Theory Ramifications 13 (2004) 193–209
- [9] **J Hoste, P D Shanahan**, *Boundary slopes of 2-bridge links determine the crossing number*, Kobe J. Math. 24 (2007) 21–39
- [10] **K Ichihara, S Mizushima**, *Crossing number and diameter of boundary slope set of Montesinos knot*, Comm. Anal. Geom. 16 (2008) 565–589
- [11] **D Johnson, C Livingston**, *Peripherally specified homomorphisms of knot groups*, Trans. Amer. Math. Soc. 311 (1989) 135–146
- [12] **P B Kronheimer, T S Mrowka**, *Witten's conjecture and Property P*, Geom. Topol. 8 (2004) 295–310
- [13] **M Macasieb, K Petersen, R Van Luijk**, *On character varieties of two-bridge knot groups* ArXiv:math.GT/09022195
- [14] **T Mattman, G Maybrun, K Robinson**, *2-Bridge knot boundary slopes: diameter and genus*, Osaka J. Math. 45 (2008) 471–489
- [15] **T Ohtsuki**, *Ideal points and incompressible surfaces in two-bridge knot complements*, J. Math. Soc. Japan 46 (1994) 51–87
- [16] **T Ohtsuki, R Riley, M Sakuma**, *Epimorphisms between 2-bridge link groups*, from: “The Zieschang Gedenschrift, Geom. Top. Monographs”, volume 14, Geom. Top. Publ., Coventry (2008) 417–450
- [17] **R Riley**, *Nonabelian representations of 2-bridge knot groups*, Quart. J. Math. Oxford Ser. (2) 35 (1984) 191–208

- [18] **D S Silver, W Whitten**, *Knot group epimorphisms*, J. Knot Theory Ramifications 15 (2006) 153–166

Pitzer College

Loyola Marymount University

jhoste@pitzer.edu, pshanahan@lmu.edu